

Read this sheet in full before using, and keep for later reference.

Find more product info online

BUFFERIN PREMIUM

Search

Fast, Highly Effective Headache Relief (headache and fever)

<Pain reliever/fever reducer>

# BUFFERIN PREMIUM

Designated Class 2 OTC drugs

Some BUFFERIN brand products contain different active ingredients. The pain relieving/fever reducing ingredients in this product are **ibuprofen** and **acetaminophen**. When consulting a doctor, dentist, pharmacist or registered seller about this product, specify that the product you are asking about contains ibuprofen and acetaminophen.

## BUFFERIN Premium Formula Four distinct features

Analgesic ingredients that effectively relieve headaches and menstrual cramps

"Quick attack" tablets\*



Easy-on-the-stomach formulation

Tablets are small and easy to swallow

\* Indications and effects include relief from headaches, fever, etc.

## Warnings

### Do not

(Failure to comply with these instructions increases the likelihood of current symptoms worsening, side effects occurring, and other potentially serious risks)

#### 1. Do not use this product if you

- (1) Have ever had an allergic reaction to this product or any of its ingredients
- (2) Have ever had an asthma attack after using this product, any other pain reliever/fever reducer, or a cold medicine
- (3) Are under 15 years old
- (4) Are pregnant and expecting to give birth within 12 weeks

#### 2. Do not take any of the following pharmaceuticals while using this product

Other pain relievers/fever reducers, cold medicines, sedatives, motion sickness medications

#### 3. Do not drive or operate machinery while using this product (users may experience drowsiness)

#### 4. Do not consume alcohol before or after taking this product

#### 5. Do not use this product continuously for extended periods

### Seek advice

#### 1. Ask a doctor, dentist, pharmacist or registered seller before use if you are

- (1) Currently receiving treatment from a doctor or dentist

- (2) Pregnant or suspect you may be pregnant
- (3) Breast-feeding
- (4) Elderly
- (5) Have ever shown allergic symptoms when taking a drug or medicine
- (6) Have been diagnosed with any of the following conditions: heart disease; kidney disease; liver disease; systemic erythematosus; mixed connective tissue disease
- (7) Have ever had any of the following illnesses: stomach or duodenal ulcer; ulcerative colitis; Crohn's disease

**2. The following symptoms may be side effects of the product. If you experience any of the following symptoms when using the product, stop use immediately and bring this sheet to a doctor, dentist, pharmacist or registered seller for consultation.**

| Affected area                 | Symptom   |
|-------------------------------|---|
| Skin                          | Rash or redness; itchiness; bruising  |
| Digestive tract               | Nausea or vomiting; loss of appetite; stomach discomfort; stomach pain; oral inflammation; heartburn; heaviness in the stomach; gastrointestinal bleeding; abdominal pain; diarrhea; bloody stool |
| Mental state / Nervous system | Dizziness   |
| Circulatory system            | Heart palpitations  |
| Respiratory system            | Shortness of breath   |
| Other                         | Blurred vision; ringing in the ears; swelling; nosebleed; bleeding from the gums; bleeding that does not stop; bleeding; back pain; excessive drop in body temperature; lethargy                  |

The following rare but serious conditions may also occur. If you experience any of the symptoms below, seek immediate medical attention from a doctor.

| Condition  | Symptoms   |
|--|--|
| Shock (anaphylaxis)  | The following symptoms occur shortly after use: itchy skin; hives; hoarse voice; sneezing; itchy throat; difficulty breathing; heart palpitations; feeling faint   |
| Mucocutaneous ocular syndrome (Stevens-Johnson syndrome), toxic epidermal necrolysis, acute generalized exanthematous pustulosis | Persistence or sudden worsening of: severe fever; bloodshot eyes; eye mucus; lip sores; sore throat; widespread rash or skin redness; small pimples (pustules) appear on reddened skin; overall lethargy; loss of appetite |
| Drug-induced hypersensitivity syndrome   | Widespread skin redness; systemic rash; fever; lethargy; swelling of lymph nodes (in the neck, armpits, groin, etc.)   |
| Impaired liver function  | Fever; itchiness; rash; jaundice (skin and whites of the eyes appear yellowish); brown urine; overall lethargy; loss of appetite   |
| Kidney disorder  | Fever; rash; reduced urine output; swelling all over the body; overall lethargy,   |

|  |  |
|--|--|
|  | arthralgia (joint pain); diarrhea  |
| Aseptic meningitis                     | Severe headache accompanied by neck stiffness; fever; nausea or vomiting (these symptoms are especially common in patients receiving care for systemic erythematosus or mixed connective-tissue disease) |
| * Myocardial infarction (heart attack) | Pressure-like chest pain; shortness of breath; cold sweat  |
| * Cerebrovascular disease (stroke)     | Sudden onset of decreased or loss of consciousness; weakness moving one side of the body; headache; vomiting, dizziness; difficulty speaking   |
| Interstitial pneumonia                 | Sudden onset or persistence of: shortness of breath or difficulty breathing after mild exertion, such as when climbing stairs; dry coughing; fever   |
| Asthma                                 | Wheezing or gasping for breath, difficulty breathing   |
| Aplastic anemia                        | Bruising; nosebleed; bleeding from the gums; fever; skin and mucous membranes appear pale; fatigue; heart palpitations; shortness of breath; feeling sick and dizziness; bloody urine                    |
| Agranulocytosis                        | Sudden onset of fever, chills or sore throat   |

**3. The following symptoms may occur during use. If these symptoms persist or worsen, stop use and bring this sheet to a doctor, pharmacist or registered seller for consultation.**

Constipation; drowsiness

**4. If symptoms do not improve after 5 or 6 doses, stop use and bring this sheet to a doctor, dentist, pharmacist or registered seller for consultation.**

### Effects

(1) Pain relieving effect on headache, pain due to stiff shoulders, menstrual pain, back pain, joint pain, neuralgia, muscle pain, sore throat, toothache, pain after tooth extraction, pain from bruising, pain from sprains, pain from bone fractures, pain from trauma and earache. (2) Reducing effect on chills and fevers

### Directions and Dosage

Avoid taking on an empty stomach and wait at least four hours between doses.

Take the following amount with lukewarm or cold water.

| Age               | Adults (15 years and up) | Children under 15 years |
|-------------------|--------------------------|-------------------------|
| Amount per 1 dose | 2 tablets                | Do not use              |
| Doses per day     | No more than 3           | ✘                       |

### Usage and Dosage Warnings

- (1) Use only as directed.
- (2) To remove the tablets from the package: As shown in the illustration on the right, press firmly with a fingertip on the protruding bubble in the clear plastic sheet that contains the tablet, pushing the tablet through the foil backing and out (swallowing the product without first removing it from the package can lead to injury, such as damage to the esophageal membrane).



### Ingredients (per 2 tablets)

| Active ingredient                 | Content | Purpose   |
|-----------------------------------|---------|---|
| Ibuprofen                         | 130 mg  | Reduces pain and fever                                      |
| Acetaminophen                     | 130 mg  |   |
| Anhydrous caffeine                | 80 mg   | Boosts pain relief effect, eases headache                   |
| Allylisopropyl acetylurea         | 60 mg   |   |
| Dehydrated aluminum hydroxide gel | 70 mg   | Protects the gastric mucosa, helps prevent stomach problems |

Inactive ingredients: cellulose, hydroxypropylcellulose, lactic acid, D-mannitol, potassium dihydrogenphosphate, silicon dioxide, magnesium stearate, polyvinyl alcohol (partially saponified), talc, titanium oxide, glycerol fatty acid ester and sodium lauryl sulfate.

### Storage and Handling

- (1) Store out of direct sunlight in a cool, dry place.
- (2) Keep out of reach of children.
- (3) Do not transfer to another container or package (may lead to incorrect use or affect product quality).
- (4) Do not use after the use-by date indicated on the package.
- (5) Avoid damaging the foil protecting tablets you are not planning to use immediately, as exposure can affect product quality.

### Inquiries

Please direct inquiries to the store where you purchased the product or the service center below.

Lion Corporation Consumer Service Center [0120-813-752](tel:0120-813-752)

Hours available: 9:00–17:00 (excluding Saturdays, Sundays and holidays)

## Lion Corporation

1-3-28 Kuramae, Taito-ku, Tokyo 111-8644

0625